

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 October 2004 (14.10.2004)

PCT

(10) International Publication Number
WO 2004/086868 A1

- (51) International Patent Classification⁷: A01N 63/02, C12N 15/82
- (21) International Application Number: PCT/GB2004/000901
- (22) International Filing Date: 3 March 2004 (03.03.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0307871.4 4 April 2003 (04.04.2003) GB
0319372.9 18 August 2003 (18.08.2003) GB
- (71) Applicant (for all designated States except US): SYNGENTA LIMITED [GB/GB]; European Regional Centre, Priestley Road, Surrey Research Park, Guildford, Surrey GU2 7YH (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CLOUGH, Martin, Stephen [GB/GB]; Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berks RG42 6EY (GB). McCAFFERY, Alan, Roy [GB/GB]; Syngenta Limited, Jealott's Hill International Research Centre, Bracknell, Berks RG42 6EY (GB).
- (74) Agents: GAAL, Jozsef, Christopher et al.; Intellectual Property Department, Syngenta Limited, Jealott's Hill International Research Centre, P.O. Box 3538, Bracknell, Berks RG42 6YA (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IMPROVED METHOD OF RESISTANCE MANAGEMENT FOR TRANSGENIC CROPS

(57) Abstract: The present invention provides a method for reducing the incidence of resistance of pests to pesticidal plants. In particular, there is provided a locus at which plant pests feed comprising at least two regions, characterised in that: a) a first region comprises plants which produce at least a first pesticidal toxin; and b) a second region comprises plants which produce at least a second pesticidal toxin; wherein a pest which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin but not the first toxin. In one aspect of the invention, the plant pests are insects. The invention also provides a method for controlling insects.

WO 2004/086868 A1

IMPROVED METHOD OF RESISTANCE MANAGEMENT FOR TRANSGENIC
CROPS

5 The present invention relates to a method for preventing or reducing the incidence of pest resistance to pesticidal plants. In particular, it relates to preventing resistance of insects to transgenic insecticidal plants such as cotton from being spread through an insect population.

10 Millions of hectares of crops worldwide are damaged each year as a result of attack by insect pests. Controlling insect pests is a serious problem for farmers who look to minimise such crop damage and resulting yield losses. This is especially true in cotton, a crop of great commercial importance. Thousands of hectares of cotton crops are damaged by a wide range of insect pests each year.

15 Entomologists calculate that crop damage caused by insects has doubled in the last 50 years, related to intensified farming efforts to feed a growing world population. The agrochemical industry has tried to control this problem with chemical solutions and insecticide spraying has become a commonplace method adopted to minimise the damage to crops. Today there are over 200 different active ingredients, in some 40,000
20 commercial chemical products, all targeted at reducing insect damage.

During one growing season, many insecticide sprays may be applied to a single crop. This intensive use of chemical insecticides imposes a high selection pressure and without careful management can lead to the rapid build-up of resistance. Examples of insect
25 resistance to pesticides have been documented worldwide. It is estimated that more than 500 arthropod pests worldwide have developed resistance to chemicals. In China, cotton yields fell by one-third between 1991 and 1993, largely due to the cotton bollworm (*Helicoverpa armigera*) which developed resistance to all the chemicals which were used for its control.

30

Insects have an exceptional ability to adapt to their environment. They have many mechanisms which allow the rapid build-up of resistance in an insect population, such as short life cycles, a high reproductive rate and the ability to travel long distances. Natural

-2 JC20 Rec'd PCT/PTO 04 OCT 2005

selection allows insects with resistance genes to survive, and the resistance trait is passed on to their offspring. Resistant insects continue to multiply as susceptible insects are eliminated by the pesticide, until eventually, insects that have a resistance gene(s) are predominant in the population and the pesticide is no longer effective. The speed with which resistance develops depends on many factors such as the rate of insect reproduction, the migration and host range of the insect, the persistence of the pesticide, the fitness costs of resistance and how often the pesticide is applied.

There are a number of known mechanisms of resistance in insects. For example, resistant insects may detoxify the toxin or remove it from their bodies faster than susceptible insects (metabolic resistance); the site where the toxin usually binds in the insect may be modified to reduce this interaction (altered target-site resistance); resistant insects may absorb the toxin slower than susceptible insects (penetration resistance); or resistant insects may detect and avoid the toxin (behavioural resistance). Pests often use more than one of these mechanisms at the same time.

One solution aimed at reducing the number of insecticide sprays and managing insect resistance is to engineer crop plants to synthesise their own insecticide. Plants may be engineered to contain, for example, insecticidal genes from other organisms. Currently the most economically significant insecticidal transgenic plants are those which contain genes from the bacterium *Bacillus thuringiensis* that produce proteins that control Lepidopteran or Coleopteran pests. Not only can the development of transgenic crops reduce the use of broad-spectrum insecticides, but they also are more target-pest specific. This means that populations of beneficial insects may not be affected. Pest control is easier with transgenic crops because, using tissue specific promoters, the insecticidal toxin can be targeted to different parts of the plant, such as the roots, which are hard to spray with conventional pesticides.

However as transgenic plants may provide extensive and continuous selection pressure on pest populations, there can be a greater potential for resistance development than with conventional insecticides. Also, unlike chemical insecticides, there are still very few genes and proteins that are known to be effective for the protection of transgenic crops against insects. In fact, only a few transgenic crops have been commercialised to date,

- 3 -

between them comprising only a handful of insecticidal genes. Therefore prevention and management of resistance build-up in populations of target insects is vitally important.

5 Insect Resistance Management (IRM) programs are designed to control such a build-up of resistance, and include the use of synthetic insecticides, biological insecticides, transgenic plants, beneficial insects, cultural practices, crop rotation, pest-resistant crop varieties and chemical attractants or deterrents.

10 One strategy employed in IRM programs is the use of refuges (refugia). In practice, refugia are often used in combination with a high dose of insecticidal agent, as currently mandated by some regulatory authorities. For transgenic crops, a refuge is the designation of a percentage of the cropped area as non-transgenic, with or without selected control treatments. Refuge areas may be within fields of transgenic or treated crops, around the border of such fields or even in adjacent fields depending on the
15 biology of the target pests. Refugia assume and work best in situations where resistance is a recessive trait. Refugia serve to maintain a population of susceptible pests. When members of this susceptible population mate with any resistant insects that emerge from protected fields, their susceptible genes dilute any resistant genes in the overall population.

20

In essence, the use of refugia serves as a mechanism for producing a population of insect pests, pests which the growers are trying to control in their field. Resistant insects which survive the insecticidal effects of feeding on the transgenic crop are not killed.

25 Prior art in the field of insect resistance management for transgenic crops relates mainly to the use of a non-transgenic refuge surrounding a transgenic crop for reducing the incidence of resistance. For example, Tang *et al.* (2001) studied the resistance of *Plutella xylostella* when feeding on transgenic broccoli expressing Cry1Ac, and demonstrated that resistance to Cry1Ac could be delayed by increasing the proportion of non-transgenic
30 refuge plants.

Roush (1997) describes four possible strategies for managing resistance to Bt-transgenic crops: (1) refuges of non-transgenic host plants preferably combined with high toxin

expression in the crop plants, (2) moderate expression of toxin in crop plants to allow susceptible insects to survive, (3) use of different toxins deployed in different varieties in a mosaic in the same area, and (4) use of varieties where each plant has a mixture of toxins. Roush describes that mosaics are the worst way to deploy two toxicants, and that
5 the most promising long term solution to resistance management is using 'pyramided' varieties containing two or more toxins. This differs from the present invention which does not include the use of mosaics. Further, although the present invention does not exclude the use of plants producing more than one insecticidal toxin, the invention *per se* does not relate to the use of pyramided varieties. In contrast to Roush, the present
10 invention teaches that the use of two regions, each comprising plants producing different insecticidal toxins, is an effective tool in insect resistance management.

Driver *et al.* (US 5,640,804) relates to the use of transgenic pest trap plants adjacent to agronomically important non-transgenic crop plants, such as walnut. The pest trap plants
15 are a preferred host for the pests, and the underlying concept is to protect crop plants by attracting pest insects away from the crop plants, and further to provide an insecticidal toxin to kill the pest insects. This is different to the present invention wherein both plant types are transgenic, neither intended as a preferred pest host, with the aim of reducing the incidence of resistance.

20 Romano (US 6,501,009) relates to the use of Cry3B and variants of Cry3B for controlling coleopteran pests. One aspect of this publication relates to an improved method of delaying the onset of resistance to Cry3B. This is achieved by using an optimised DNA cassette to achieve higher levels of expression of Cry3B and/or by
25 simultaneously exposing target insects to both Cry3 toxins and other insecticidal proteins. Further, 'simultaneous exposure' refers to the expression of Cry3B and another insecticidal protein in the same plant using a gene stacking type approach. This is different to the present invention, where different insecticidal toxins are produced in different plants.

30 At its best, the current refuge strategy works well, but this may not be the case in all circumstances. For example, a refuge strategy works well when the resistance trait to be controlled is a recessive trait. However, if the resistance trait is dominant, the refuge will

be much less effective at controlling resistance, and at best will only slowly dilute out the resistance trait. Also, refugia work best for IRM when combined with a crop expressing a high dose of insecticidal toxin so that all insects which are heterozygous for a resistance allele are killed. This is to ensure that the resistance trait is recessive in nature.

- 5 However, it is often difficult to ensure expression of a sufficiently high enough dose of toxin to kill insects which are heterozygous for the resistance allele, and often difficult to ensure such high doses are expressed all season long.

The present patent application describes an improved method for IRM, which works to
10 control recessive or dominant resistance traits better than using a refuge. This improved method may also be used in conjunction with a refuge. The invention uses a region comprising plants which produce at least one insecticidal toxin (principal crop plants), said toxin being different to the toxin produced by the plants in a second region. The effect of the different insecticidal toxins is to kill insects which are resistant to the toxin
15 of the principal crop plants rather than allowing them to breed and thus spread the resistance trait. In this way insects are unlikely to survive as this would require resistance to at least two different insecticidal toxins preferably having different modes of action. Therefore the incidence of resistance is reduced. This invention may be used in conjunction with other IRM techniques.

20 According to the present invention there is provided a locus at which plant pests feed comprising at least two regions, characterised in that: a) a first region comprises plants which produce at least a first pesticidal toxin; and b) a second region comprises plants which produce at least a second pesticidal toxin; wherein a pest which can develop
25 resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin but not the first toxin. In one embodiment of the invention, the plant pests are selected from the group consisting of insects, mites and nematodes. In further embodiment of the invention, the plant pests are
30 insects. In this aspect of the invention, the plants produce insecticidal toxins.

The word insecticidal as used herein describes the effect of a toxin on insects. It is not limited to death of the insect, but also includes any effect which is detrimental to the

insect, for example sickness, anti-feedant activity, growth retardation, reduced reproductive ability and reduced fecundity.

Resistant insects are those that do not suffer any substantial or appreciable detrimental effects as a result of exposure to or ingestion of a suitable dose of insecticidal toxin. A suitable dose of insecticidal toxin may be measured by exposure to or ingestion of the toxin by a susceptible insect and identification of the dose at which detrimental effect(s) are observed. Detrimental effects to the insect are described above in the definition of the word insecticidal. These detrimental effects will reduce the incidence of transfer of a resistance trait from a resistant insect to future generations of insects. In one aspect of the invention, the resistance trait is a dominant trait.

The word plants as used herein refers to plants and plant parts and includes seeds.

The invention includes plants which produce more than one toxin, for example via gene stacking. The plants of either the first and/or second region may even produce the same toxins, with the *proviso* that the first region comprises plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin but not the first toxin. The invention is not limited to loci which comprise first and second regions that only comprise plants which produce insecticidal toxin(s), but may also contain other plants in addition. The other plants may be non-transgenic. In one aspect of the invention, the plants of either region may also produce toxins to make them resistant to non-insect pests such as viruses, fungi or nematodes. In another aspect of the invention, the plants of either region may be tolerant to chemical herbicides. In a further aspect of the invention, the locus may comprise more than two regions, wherein said additional regions may comprise plants which produce insecticidal toxins. In one embodiment of the invention, the locus may comprise a third region, which region comprises non-insecticidal plants.

The skilled man will be familiar with insects which feed at the locus. Preferably, insects which feed at the locus include pest insects which cause damage to plants. More preferably, this includes insects which are, or can develop to be, resistant to an insecticidal toxin. More preferably still, it includes insects selected from the group

- comprising: *Acanthoscelides obtectus*, *Bruchus* spp., *Callosobruchus* spp. (bruchid beetles), *Agriotes* spp. (wireworms), *Amphimallon* spp. (chafer beetles), *Anthonomus grandis* (cotton boll weevil), *Ceutorhynchus assimilis* (cabbage seed weevil), *Cylas* spp. (sweet potato weevils), *Diabrotica* spp. (corn root worms), *Epicauta* spp. (black blister beetles), *Epilachna* spp. (melon beetles etc.), *Leptinotarsa decemlineata* (Colorado potato beetle) *Meligethes* spp. (blossom beetles), *Melolontha* spp. (cockchafer), *Phyllotreta* spp., *Psylliodes* spp. (flea beetles), *Popillia japonica* (Japanese beetle), *Scolytus* spp. (bark beetles), *Acleris* spp. (fruit tree tortrix), *Acraea acerata* (sweet potato butterfly), *Agrotis* spp. (cutworms), *Autographa gamma* (silver-Y moth), *Chilo* spp. (stalk borers), *Cydia pomonella* (codling moth), *Diparopsis* spp. (red bollworms), *Ephestia* spp. (warehouse moths), *Heliothis* spp., *Helicoverpa* spp. (budworms, bollworms), *Mamestra brassicae* (cabbage moth), *Manduca* spp. (hornworms), *Maruca testulalis* (mung moth), *Mythimna* spp. (cereal armyworms), *Ostrinia nubilalis* (European corn borer), *Pectinophora gossypiella* (pink bollworm), *Phthorimaea operculella* (potato tuber moth), *Pieris brassicae* (large white butterfly), *Pieris rapae* (small white butterfly), *Plodia interpunctella* (Indian grain moth), *Plutella xylostella* (diamond-back moth), *Pseudoplusia includens* (soybean looper), *Sitotroga cerealella* (Angoumois grain moth), *Spodoptera* spp. (armyworms), *Trichoplusia ni* (cabbage semilooper), *Acheta* spp. (field crickets), *Grylotalpa* spp. (mole crickets), *Locusta migratoria* (migratory locust), *Schistocerca gregaria* (desert locust), *Acrythosiphon pisum*, *Drosophila* spp., *Acrosternum hilare* (green stink bug), *Aphis gossypii* (cotton aphid), *Campylomma liebenehti* (apple dimpling bug), *Creontiades dilutus* (green mirid), *Crociosema plebejana* (cotton tipworm), *Earias huegelli* (rough bollworm), *Euschistus servus* (brown stink bug), *Frankliniella* spp. (thrips), *Lygus lineolaris* (tarnished plant bug), *Tetranychus urticae* (spider mite) and *Thrips tabaci* (onion thrips).

Most preferably, it includes insects selected from the group comprising: *Anthonomus grandis* (cotton boll weevil), *Pectinophora gossypiella* (pink bollworm), *Heliothis virescens* (tobacco budworm), *Helicoverpa zea* (cotton bollworm), *Helicoverpa armigera* (American bollworm), *Helicoverpa punctigera* (native bollworm), *Spodoptera exigua* (beet armyworm) and *Spodoptera frugiperda* (fall armyworm).

With the benefit of the present disclosure, the skilled man will be familiar with insecticidal toxins that can be expressed in plants which may be suitable for use in this invention. Suitable toxins may even be those known in the prior art. For example, they include crystal proteins from *Bacillus thuringiensis*, many of which have been

5 extensively studied and are well known in the prior art such as Cry1Ac, Cry2Ab and Cry1F. A non-limiting list of protein toxins from *Bacillus thuringiensis* which may be used in the present invention is available on the internet at http://www.biols.susx.ac.uk/home/Neil_Crickmore/Bt/index.html. Further non-limiting examples of insecticidal toxins are vegetative insecticidal proteins VIP3A and VIP3B

10 from *Bacillus thuringiensis*, 445 from *Paecilomyces farinosus* (see International Patent Application publication number WO01/00841) and GGK (see International Patent Application publication number WO02/098911). Alternative suitable insecticidal toxins may, for example, be isolated from bacteria, fungi, plants or other sources. The genes encoding these toxins can be cloned and transformed into suitable plants under the

15 control of a plant-operable gene cassette, using standard molecular and cell biology techniques. The toxins may be targeted to particular parts of the plant such as the roots, leaves or seeds by cloning the genes encoding the toxins to be under the control of tissue-specific promoters. Alternatively, the toxins may only be produced at a certain growth stage of the plant through use of inducible or temporal promoters. The first and second

20 insecticidal toxins may be insecticidal towards different spectra of insect species. Preferably the first and second toxins are insecticidal towards the same or similar insect species, or overlapping spectra of insect species. Preferably the first and second toxins act at different binding sites to one another. More preferably, the first and second toxins have different modes of action to one another.

25

In one aspect of the invention, the plants of the first and second regions may optionally exhibit other beneficial traits, which also may have been introduced via gene cloning and plant transformation. Any number of these traits may be stacked with the insecticidal toxin in the plants. For example, the plants of either region may exhibit resistance to a

30 particular herbicide, fungal disease, viral infection or nematode infestation. An example of resistance to a herbicide is described in International Patent Application publication number WO 00/66747 wherein a mutant form of the enzyme EPSP synthase is expressed in a plant so that the plant is tolerant to the herbicide glyphosate.

In a particular embodiment of the present invention, the second region is within a mile from the first region. In another embodiment of the present invention, the second region is within a quarter of a mile from the first region. In a further embodiment the second region is adjacent to the first region. In a still further embodiment the second region is a border around the perimeter of the first region. In a further embodiment the second region comprises one or more strips within the first region. In a further embodiment the locus comprises a random distribution of first and second regions within the locus. The schematic diagrams provided in Figures 1 to 10 represent non-limiting examples of the possible arrangement of first and second regions within the locus of the present invention. In one aspect of the invention, a mosaic pattern of first and second regions may be specifically excluded. In an embodiment where the locus comprises a third region, which region comprises non-insecticidal plants, said third region may also be arranged with respect to the first and second regions according to the arrangements described above.

15 A locus according to the present invention may be comprised by a farm, wherein the at least two regions are fields. The following non-limiting examples describe possible arrangements of the fields. For example, the fields may be adjacent to one another (see Figures 5 to 7). Alternatively, the locus of the present invention may be a field, wherein the at least two regions are areas of the field comprising different plants. The second region may be arranged, for example, as a border around the perimeter of the first region (see Figure 1), as a series of horizontal or vertical strips amongst the first regions (see Figures 2 and 3) or as a block within the first region (see Figure 4). The deployment of the regions within the locus is dependent on the biology or mobility of the target pest. In one aspect of the invention, there may be a border around the locus comprising or consisting of non-insecticidal plants, non-transgenic plants, non-host plants or being uncultivated. In another aspect of the invention, the locus may comprise a third region, said region being, for example, a border around the first and / or second regions, and comprising non-insecticidal plants.

30 Alternatively, a locus according to the present invention may be comprised by, for example, a garden, forest, glasshouse or seed store. The locus may even be comprised by a lake such that the invention is used to control aquatic insects. However, the scope of

this invention is clearly restricted to loci wherein the invention would be functional. The skilled man would understand that the present invention excludes the possibility of the locus being the world, wherein the first region is America and the second region is Europe, for example.

5 In a particular embodiment of the present invention the locus comprises at least two regions wherein the first region comprises plants which produce at least a first insecticidal toxin, and the second region comprises plants which produce at least a second insecticidal toxin, wherein the first insecticidal toxin has a different mode of
10 action to the second insecticidal toxin. Examples of known modes of action of insecticidal chemicals and toxins include, but are not limited to acetyl choline esterase inhibitors, GABA-gated chloride channel antagonists, sodium channel modulators, acetyl choline receptor modulators, chloride channel activators, juvenile hormone mimics, fumigants, selective feeding blockers, growth inhibitors, disrupters of insect midgut
15 membranes, inhibitors of oxidative phosphorylation, disrupters of ATP formation, oxidative phosphorylation uncouplers, inhibitors of magnesium stimulated ATPase, inhibitors of chitin biosynthesis, ecdysone agonist or disrupters, electron transport inhibitors and voltage dependant sodium channel blockers. For example, the first toxin may be a disrupter of the insect midgut membrane, such as a crystal toxin from *Bacillus*
20 *thuringiensis*, and the second toxin may be a growth inhibitor.

In a further embodiment of the invention, the first insecticidal toxin is a crystal protein from *Bacillus thuringiensis* and the second insecticidal toxin is a vegetative insecticidal protein (VIP) from *Bacillus thuringiensis*, or vice versa. Many crystal proteins from
25 *Bacillus thuringiensis* have been isolated and are known to have an insecticidal effect. Preferably, the crystal protein of the present invention is Cry1Ac. Preferably the VIP protein is VIP3A. In one aspect of the invention, the plants which produce either the first or second insecticidal toxin may comprise both Cry1Ac and Cry2Ab.

30 In a further embodiment of the present invention, the plants which comprise the first toxin and the plants which comprise the second toxin are from different genera. For example, the plants which comprise the first toxin may be cotton plants from the genus *Gossypium* L., and the plants which comprise the second toxin may be corn plants from

the genus *Zea* L.. In further non-limiting example, the plants which comprise the first toxin may be wheat plants from the genus *Triticum* L. and the plants which comprise the second toxin may be barley plants from the genus *Hordeum* L..

- 5 In a further embodiment of the present invention, the plants which comprise the first toxin and the plants which comprise the second toxin are from the same genus. In a further embodiment of the present invention, the plants which comprise the first toxin and the plants which comprise the second toxin are cotton plants. Preferably the plants are of the same species. More preferably the plants are the Upland Cotton species,
10 *Gossypium hirsutum*. The cotton plants of the first and second regions may be the same or different varieties.

- In a further embodiment of the invention, at least 5% of the locus comprises the first region and at least 5% of the locus comprises the second region. In one alternative, at
15 least 20% of the locus comprises the first region and at least 20% of the locus comprises the second region. In another alternative, 50% of the locus comprises the first region and 50% of the locus comprises the second region.

- Further, the invention provides the option of applying a chemical spray to some or all of
20 the regions or parts of said regions within the locus. The chemical may, for example, be an insecticide, fungicide or herbicide. Preferably such an insecticidal chemical acts at a different binding site to the insecticidal toxins produced by the plants of the first and / or second regions. Preferably such an insecticidal chemical has a different mode of action to the insecticidal toxins produced by the plants of the first and / or second regions.
25 More preferably, the chemical is not a *Bacillus thuringiensis* insecticide.

- According to the present invention there is provided a method of controlling insects comprising providing a locus at which insects feed comprising at least two regions, characterised in that: a) a first region comprises plants which produce at least a first
30 insecticidal toxin; and b) a second region comprises plants which produce at least a second insecticidal toxin; wherein an insect which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises

plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin but not the first toxin.

The words 'controlling' or 'control' as used herein refer not just to death of insects, but also include other detrimental effects on insects such as sickness, anti-feedant activity, growth retardation, reduced reproductive ability and reduced fecundity.

In an embodiment of the invention, a locus as described above is used in a method of controlling insects.

10

According to the present invention there is provided a method of reducing the incidence of resistance to a first insecticidal toxin comprising the steps of providing a locus at which insects feed comprising at least two regions, characterised in that: a) a first region comprises plants which produce at least a first insecticidal toxin; and b) a second region comprises plants which produce at least a second insecticidal toxin; wherein an insect which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second when the plants of the second region produce the second toxin but not the first, so that insects which have developed or are developing resistance to the first insecticidal toxin are controlled by the second toxin.

20

The terms 'developed resistance' and 'developing resistance' refer to resistance within a population of insects rather than individual insects. While it may be possible for an individual insect to apparently become resistant to an insecticidal toxin, for example by the overproduction of a detoxification enzyme in response to ingestion of the toxin, this ability to induce enzyme production is likely pre-determined as a result of a mutation in the insect genome. Therefore, insects hatch as either resistant or susceptible to a particular toxin. The terms 'developed resistance' or 'developing resistance' encompass the development of resistance via evolution through generations of breeding.

25
30

In an embodiment of the invention, a locus as described above is used in a method of reducing the incidence of resistance of insects to a first insecticidal toxin.

In a further embodiment of the present invention, there is a method as described above or a locus as described above wherein either the first or second region comprises Bollgard® cotton plants. Preferably either the first or second region comprises Bollgard I® cotton plants which produce the insecticidal toxin Cry1Ac. More preferably either the first or second region comprises Bollgard II® cotton plants which produce the insecticidal toxins Cry1Ac and Cry2Ab in the same plant.

In a further embodiment of the present invention, there is a method as described above or a locus as described above wherein the first region comprises Bollgard® cotton plants and the second region comprises VIP cotton plants. In a further embodiment still, there is a method as described above or a locus as described above wherein the first region comprises VIP cotton plants and the second region comprises Bollgard® cotton plants. The Bollgard® cotton plants may be Bollgard I® or Bollgard II®. The VIP cotton plants produce the insecticidal toxin VIP3A (see, for example, International Patent Application Number PCT/EP03/11725). In a further aspect of the invention, the first region comprises VIP cotton plants and the second region comprises Bollgard® cotton plants.

In a further embodiment of the present invention, there is provided a method or a locus as described above, wherein the first or second region comprises plants expressing the insecticidal toxin Cry3A or a modified version thereof. In a further embodiment of the present invention, there is provided a method or a locus as described above, wherein the first region comprises plants which comprise Cry3A toxin from *Bacillus thuringiensis* or a modified version thereof, and the second region comprises plants which comprise Cry3B toxin from *Bacillus thuringiensis*. Modified versions of Cry3A include, for example, Cry3A toxins comprising a cathepsin G protease recognition site in domain 1, such as those described in International Patent Application publication number WO03/18810.

In addition to loci at which insects feed comprising plants expressing insecticidal toxins for reducing the incidence of resistance in insects, the present invention further includes loci at which nematodes or mites feed, comprising plants expressing nematocidal or mitocidal proteins for reducing the incidence of resistance in nematodes or mites respectively, and methods and uses thereof.

In one aspect of the present invention neither the plants of the first region, nor the plants of the second region produce a Cry3 toxin. In a further aspect of the present invention neither the plants of the first region, nor the plants of the second region produce a Cry3B toxin. For example, the present invention may specifically exclude a locus wherein one of the regions of said locus comprises plants which produce a Cry3B toxin or variants thereof.

DESCRIPTION OF THE FIGURES

10 The accompanying figures illustrate non-limiting examples of the arrangement of the first and second regions within the locus.

Figure 1 – Second region forms a border around the perimeter of the first region

Figure 2 – Second region forms brackets either side of the first region

15 Figure 3 – Second region forms a series of horizontal or vertical strips amongst the first regions

Figure 4 – Second region forms a block within the first region

Figures 5 – 7 – First and second regions are adjacent

Figure 8 – 10 – A plurality of first and second regions

20

REFERENCES

Driver J. and Dandekar A.M. (1997) Pest trap plants and crop protection. US 5,640,804.

25 Romano C.P. (2002) Expression of Cry3B insecticidal protein in plants. US 6,501,009.

Roush R.T. (1997) Bt-transgenic crops: just another pretty insecticide or a chance for a new start in resistance management. *Pesticide science* 51(3), 328-334.

30 Tang J.D., Collins H.L., Metz T.D., Earle E.D., Zhao J.Z., Roush R.T. and Shelton A.M. (2001) Greenhouse tests on resistance management of Bt transgenic plants using refuge strategies. *Journal of Economic Entomology* 94(1), 240-247.

CLAIMS

1. A locus at which plant pests feed comprising at least two regions, characterised in that:
 - 5 a) a first region comprises plants which produce at least a first pesticidal toxin; and
 - b) a second region comprises plants which produce at least a second pesticidal toxin; wherein a pest which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin
- 10 but not the first toxin.
2. A locus according to claim 1, wherein the plant pests are selected from the group consisting of insects, mites and nematodes.
- 15 3. A locus according to claim 2, wherein the plant pests are insects.
4. A locus according to claim 3 also comprising a third region, which region comprises non-insecticidal plants.
- 20 5. A locus according to any one of claims 1 to 4, wherein the second region is within a mile from the first region.
6. A locus according to claim 5, wherein the second region is within a quarter of a mile from the first region.
- 25 7. A locus according to any of claims 1 to 6, wherein the second region is adjacent to the first region.
8. A locus according to any of claims 1 to 6, wherein the second region is a border
- 30 around the perimeter of the first region.
9. A locus according to any of claims 1 to 6, wherein the second region comprises one or more strips within the first region.

- 16 -

10. A locus according to any of claims 1 to 6 which comprises a random distribution of first and second regions.

11. A locus according to any one claims 3 to 10, wherein the first insecticidal toxin
5 has a different binding site to the second insecticidal toxin.

12. A locus according to any one of claims 3 to 11, wherein the first insecticidal toxin has a different mode of action to the second insecticidal toxin.

10 13. A locus according to any of claims 3 to 10, wherein the first insecticidal toxin is a crystal protein from *Bacillus thuringiensis* and the second insecticidal toxin is a VIP protein from *Bacillus thuringiensis*, or vice versa.

14. A locus according to any of the preceding claims, wherein the plants which
15 comprise the first toxin and the plants which comprise the second toxin are from different genera.

15. A locus according to any of the preceding claims, wherein the plants which
comprise the first toxin and the plants which comprise the second toxin are from the
20 same genus.

16. A locus according to claim 15, wherein the plants which comprise the first toxin and the plants which comprise the second toxin are cotton plants.

25 17. A locus according to any of the preceding claims, wherein at least 5% of the locus comprises the first region and least 5% of the locus comprises the second region.

18. A locus according to claim 17, wherein at least 20% of the locus comprises the first region and at least 20% of the locus comprises the second region.

30

19. A locus according to claim 18, wherein 50% of the locus comprises the first region and 50% of the locus comprises the second region.

20. A method of controlling insects comprising providing a locus at which insects feed comprising at least two regions, characterised in that:

- a) a first region comprises plants which produce at least a first insecticidal toxin; and
- b) a second region comprises plants which produce at least a second insecticidal toxin;

wherein an insect which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second toxin when the plants of the second region produce the second toxin but not the first toxin.

21. Use of a locus according to any of claims 3 to 19 in a method of controlling insects.

22. A method of reducing the incidence of resistance of an insect to a first insecticidal toxin comprising the steps of providing a locus at which insects feed comprising at least two regions, characterised in that:

- a) a first region comprises plants which produce at least a first insecticidal toxin; and
- b) a second region comprises plants which produce at least a second insecticidal toxin;

wherein an insect which can develop resistance to the first toxin does not develop resistance to the second toxin, and the first region comprises plants which produce the first toxin but not the second when the plants of the second region produce the second toxin but not the first, so that insects which have developed or are developing resistance to the first insecticidal toxin are controlled by the second toxin.

23. Use of a locus according to any of claims 3 to 19 in a method of reducing the incidence of resistance of insects to a first insecticidal toxin.


24. A method according to claim 20 or 22 or a locus according to any of claims 3 to 19, wherein either the first or second region comprises Bollgard® cotton plants.

- 18 -

25. A method according to claim 20 or 22 or a locus according to any of claims 3 to 19, wherein the first region comprises Bollgard® cotton plants and the second region comprises VIP cotton plants, or vice versa.
- 5 26. A method according to claim 20 or 22 or a locus according to any of claims 3 to 10, wherein the first region comprises plants which comprise Cry3A toxin from *Bacillus thuringiensis* or a modified version thereof, and the second region comprises plants which comprise Cry3B toxin from *Bacillus thuringiensis*.


1/5

FIGURE 1



□ FIRST REGION
▨ SECOND REGION

FIGURE 2



□ FIRST REGION
▨ SECOND REGION

2 / 5 **JC20 Rec'd PCT/PTO 04 OCT 2005**

FIGURE 3





FIGURE 4



3 / 5

FIGURE 5





FIGURE 6



4 / 5

FIGURE 7






FIGURE 8




5/5

FIGURE 9



□ FIRST REGION
 ▨ SECOND REGION

FIGURE 10



□ FIRST REGION
 ▨ SECOND REGION

INTERNATIONAL SEARCH REPORT

GB2004/000901

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01N63/02 C12N15/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | WHALON, M.E., ET AL.: "Chapter 7: Bacillus thuringiensis: use and resistance management" INSECTICIDAL WITH NOVEL MODES OF ACTION. MECHANISM AND APPLICATION. SPRINGER, BERLIN. EDS ICHAAAYA, I. ET AL., 1998, pages 106-137, XP001196940 page 122 - page 124 | 1-26 |
| X | WO 02/13609 A (AROA IN RAFFI V ; UNIV CALIFORNIA (US)) 21 February 2002 (2002-02-21) claims 7,8,12 ----- -/- | 1-26 |

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

1 July 2004

Date of mailing of the international search report

29/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

GB2004/000901

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|--|--|---------------------------------------|
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | ROUSH R T: "Two-toxin strategies for management of insecticidal transgenic crops: Can pyramiding succeed where pesticide mixtures have not?" PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY OF LONDON B BIOLOGICAL SCIENCES, vol. 353, no. 1376, 29 October 1998 (1998-10-29), pages 1777-1786, XP008032212 ISSN: 0962-8436 abstract section 5a and 5b page 1780 | 1-26 |
| X | FRUTOS ROGER ET AL: "Managing insect resistance to plants producing <i>Bacillus thuringiensis</i> toxins" CRITICAL REVIEWS IN BIOTECHNOLOGY, vol. 19, no. 3, September 1999 (1999-09), pages 227-276, XP008032140 ISSN: 0738-8551 page 256 section C | 1-26 |
| X | " <i>Bacillus thuringiensis</i> Cry2Ab2 protein and the genetic material necessary for its production in cotton (006487) fact sheet" US ENVIRONMENTAL PROTECTION AGENCY. BIOPESTICIDE ACTIVE INGREDIENT FACT SHEET., 'Online! XP002286661 23-12-2002 Retrieved from the Internet: URL: http://www.epa.gov/opppbpd1/biopesticides/ingredients/factsheets/factsheet_006487.htm the whole document | 1-26 |
| X | EP 0 400 246 A (PLANT GENETIC SYSTEMS NV) 5 December 1990 (1990-12-05) the whole document | 1-26 |
| P,X | WO 03/060463 A (MURDOCK LARRY LEE ; GAFFNEY PATRICK JOSEPH (US); PITTENDRIGH BARRY ROB) 24 July 2003 (2003-07-24) page 34, line 4 - page 35, line 6; claims 78-114; figures 8-11 | 1-3,5,6, 11,12, 14,15, 20-23 |
| A | CAPRIO MICHAEL A: "Evaluating resistance management strategies for multiple toxins in the presence of external refuges" JOURNAL OF ECONOMIC ENTOMOLOGY, vol. 91, no. 5, October 1998 (1998-10), pages 1021-1031, XP001189686 ISSN: 0022-0493 the whole document | 7-10 |
| | -/-- | |

INTERNATIONAL SEARCH REPORT

GB2004/000901

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| A | HUNT, T.E, ET AL.: "Resistance management for European corn borer and Bt transgenic corn: refuge design and placement" NEBRASKA COOPERATIVE EXTENSION. NEBFACTS. NF00-425, 'Online! October 2002 (2002-10), XP002286662 Retrieved from the Internet: URL: http://ianrpubs.unl.edu/insects/nf425.htm figures 1,2 | 7-9 |
| A | US 6 501 009 B1 (ROMANO CHARLES P) 31 December 2002 (2002-12-31) column 8, line 16 - line 39 | 13,24-26 |
| A | WO 96/08136 A (TREETECH MANAGEMENT INC DOING ; UNIV CALIFORNIA (US)) 21 March 1996 (1996-03-21) claims 1-24 | 1-26 |
| A | WO 01/92561 A (GAFFNEY PATRICK JOSEPH ; MURDOCK LARRY LEE (US); PITTENDRIGH BARRY ROB) 6 December 2001 (2001-12-06) the whole document | |

INTERNATIONAL SEARCH REPORT

Information on patent family members

GB2004/000901

| Patent document cited in search report | | Publication date | Patent family member(s) | Publication date |
|---|----|---------------------|----------------------------|---------------------|
| WO 0213609 | A | 21-02-2002 | AU 9125901 A | 25-02-2002 |
| | | | CA 2418365 A1 | 21-02-2002 |
| | | | EP 1307098 A1 | 07-05-2003 |
| | | | WO 0213609 A1 | 21-02-2002 |
| EP 0400246 | A | 05-12-1990 | EP 0400246 A1 | 05-12-1990 |
| | | | AT 109511 T | 15-08-1994 |
| | | | AU 623530 B2 | 14-05-1992 |
| | | | AU 5724590 A | 07-01-1991 |
| | | | CA 2032481 A1 | 01-12-1990 |
| | | | CA 2284255 A1 | 01-12-1990 |
| | | | DE 69011232 D1 | 08-09-1994 |
| | | | DE 69011232 T2 | 08-12-1994 |
| | | | DK 408403 T3 | 28-11-1994 |
| | | | WO 9015139 A1 | 13-12-1990 |
| | | | EP 0408403 A1 | 16-01-1991 |
| | | | ES 2060975 T3 | 01-12-1994 |
| | | | HK 1008050 A1 | 30-04-1999 |
| | | | JP 4500161 T | 16-01-1992 |
| | | | JP 3325564 B2 | 17-09-2002 |
| | | | US 6172281 B1 | 09-01-2001 |
| | | | US 5908970 A | 01-06-1999 |
| | | | US 5866784 A | 02-02-1999 |
| WO 03060463 | A | 24-07-2003 | WO 03060463 A2 | 24-07-2003 |
| US 6501009 | B1 | 31-12-2002 | US 2003115630 A1 | 19-06-2003 |
| WO 9608136 | A | 21-03-1996 | US 5640804 A | 24-06-1997 |
| | | | AT 212182 T | 15-02-2002 |
| | | | AU 3595795 A | 29-03-1996 |
| | | | CA 2219915 A1 | 21-03-1996 |
| | | | DE 69525165 D1 | 14-03-2002 |
| | | | DE 69525165 T2 | 02-10-2002 |
| | | | DK 784421 T3 | 13-05-2002 |
| | | | EP 0784421 A1 | 23-07-1997 |
| | | | ES 2170808 T3 | 16-08-2002 |
| | | | PT 784421 T | 31-07-2002 |
| | | | WO 9608136 A1 | 21-03-1996 |
| | | | ZA 9507691 A | 27-05-1996 |
| WO 0192561 | A | 06-12-2001 | AU 7521101 A | 11-12-2001 |
| | | | CA 2410584 A1 | 06-12-2001 |
| | | | EP 1287352 A2 | 05-03-2003 |
| | | | WO 0192561 A2 | 06-12-2001 |